

Correspondence between Calogero-Moser systems and integrable $SL(N, \mathbb{C})$ Euler-Arnold tops

Andrey Smirnov

E-mail: asmirnov@itep.ru

Moscow Institute of Physics and Technology, Moscow, Russia

Institute for Theoretical and Experimental Physics, Moscow, Russia

Abstract

The equivalence between the N -particle Calogero-Moser systems and the integrable $sl(N, \mathbb{C})$ -tops is shown. New rational and trigonometric classical Lax operators for these systems are found. Relations with new solutions of the classical Yang-Baxter equations for $sl(N, \mathbb{C})$ are discussed. Explicit formulae for $N=2,3,4$ are presented.

1 Introduction

It was shown in [1] that the systems of the $sl(2, \mathbb{C})$ Euler-Arnold tops [2, 3, 4] are equivalent to the two-particle Calogero-Moser systems (CM). In the present paper we generalize these results to an arbitrary rank. We show that the N -particle trigonometric and rational CM systems correspond to some new classes of $sl(N, \mathbb{C})$ tops. The L-operators for these systems do not coincide with the standard degenerations of the elliptic Belavin L-operator. Thus, these Lax operators correspond to some new solutions of the classical Yang-Baxter equation for $sl(N, \mathbb{C})$. In appendix we present explicit expressions for these L-operators for $N=2,3,4$. We write out the integrals of motion and then check that they commute with respect to the standard Poisson structure on $sl(N, \mathbb{C})$.

2 Basic example in rank one

The classical elliptic L-operator for $\mathfrak{sl}(2, \mathbb{C})$ is the solution of the classical equation for the linear Poisson bracket:

$$\{L^{ET}(z) \otimes 1, 1 \otimes L^{ET}(z)\} = [r(z), 1 \otimes L^{ET}(z) + L^{ET}(z) \otimes 1], \quad (1)$$

where $r(z)$ is the classical elliptic Baxter r-matrix for $\mathfrak{sl}(2, \mathbb{C})$. Explicitly for $L^{ET}(z)$ we get:

$$L^{ET}(z) = \begin{bmatrix} -\frac{\theta'_{11}(0) \theta_{11}(z - \frac{1}{2})}{\theta_{11}(z) \theta_{11}(-\frac{1}{2})} S_3 & \frac{\theta'_{11}(0) e^{-\pi i z}}{\theta_{11}(z)} \left(\frac{\theta_{11}(z - \frac{\tau}{2})}{\theta_{11}(-\frac{\tau}{2})} S_1 + \frac{\theta_{11}(z - \frac{1+\tau}{2})}{\theta_{11}(-\frac{1+\tau}{2})} S_2 \right) \\ \frac{\theta'_{11}(0) e^{-\pi i z}}{\theta_{11}(z)} \left(\frac{\theta_{11}(z - \frac{\tau}{2})}{\theta_{11}(-\frac{\tau}{2})} S_1 - \frac{\theta_{11}(z - \frac{1+\tau}{2})}{\theta_{11}(-\frac{1+\tau}{2})} S_2 \right) & \frac{\theta'_{11}(0) \theta_{11}(z - \frac{1}{2})}{\theta_{11}(z) \theta_{11}(-\frac{1}{2})} S_3 \end{bmatrix}.$$

Here $\theta_{k m}(z)$ are the standard theta functions with the elliptic parameter τ :

$$\theta_{k m}(z, \tau) = \sum_{n \in \mathbb{Z}} \exp \left(2 \pi i \left(\left(n + \frac{k}{2} \right)^2 \frac{\tau}{2} + \left(z + \frac{m}{2} \right) \left(n + \frac{k}{2} \right) \right) \right). \quad (2)$$

Equation (1) fixes the Poisson brackets between S_i :

$$\{S_i, S_j\} = 2i \epsilon_{ijk} S_k.$$

In what follows we also use the Chevalley basis:

$$S_2 = i(S^+ - S^-), \quad S_1 = S^+ + S^-, \quad \{S_3, S^+\} = 2S^+, \quad \{S_3, S^-\} = -2S^-, \quad \{S^+, S^-\} = S_3.$$

In [5], in the framework of Hitchin approach to the integrable systems [6], the interrelations between the elliptic $\mathfrak{sl}(2, \mathbb{C})$ top and the elliptic two particle Calogero-Moser system was found. In terms of the Lax operators for these systems the correspondence is looks like a gauge transformation:

$$L^{ET}(z) = \Xi(z) L^{ECM}(z) \Xi^{-1}(z), \quad (3)$$

where $\Xi(z)$ is a 2×2 matrix depending on the spectral parameter z and on the relative coordinate $u = u_1 - u_2$ of the CM system:

$$\Xi(z) = \Xi(z, u) = \begin{bmatrix} \theta_{00}(z - 2u, 2\tau) & -\theta_{00}(z + 2u, 2\tau) \\ -\theta_{10}(z - 2u, 2\tau) & \theta_{10}(z + 2u, 2\tau) \end{bmatrix}, \quad (4)$$

and $L^{ECM}(z)$ is the L-operator for the elliptic CM system:

$$L^{ECM} = \begin{bmatrix} v & \nu \frac{\theta_{11}(z+u) \theta'_{11}(0)}{\theta_{11}(u) \theta_{11}(z)} \\ \nu \frac{\theta_{11}(z-u) \theta'_{11}(0)}{\theta_{11}(-u) \theta_{11}(z)} & -v \end{bmatrix}.$$

From (3) we have:

$$\begin{cases} S_1 = -\frac{\theta_{10}(0) \theta_{10}(2u)}{\theta'_{11}(0) \theta_{11}(2u)} v + \frac{\theta_{10}^2(0) \theta_{00}(2u) \theta_{01}(2u)}{\theta_{00}(0) \theta_{01}(0) \theta_{11}^2(2u)} \nu, \\ S_2 = \frac{\theta_{00}(0) \theta_{00}(2u)}{i \theta'_{11}(0) \theta_{11}(2u)} v - \frac{\theta_{00}^2(0) \theta_{10}(2u) \theta_{01}(2u)}{i \theta_{10}(0) \theta_{01}(0) \theta_{11}^2(2u)} \nu, \\ S_3 = -\frac{\theta_{01}(0) \theta_{01}(2u)}{\theta'_{11}(0) \theta_{11}(2u)} v + \frac{\theta_{01}^2(0) \theta_{00}(2u) \theta_{10}(2u)}{\theta_{00}(0) \theta_{10}(0) \theta_{11}^2(2u)} \nu. \end{cases} \quad (5)$$

Our general aim is to find the trigonometric and rational analogs for these formulae. In the limit $\tau \rightarrow i\infty$ corresponding to the trigonometric degeneration expressions for S_i (5) diverge. To overcome this problem let us apply to (3) an additional gauge transformation [7, 8] depending on $q = \exp(\pi i \tau)$:

$$A^T(q) L^{ET}(z) A^T(q)^{-1} = A^T(q) \Xi(z) L^{ECM}(z) \Xi^{-1}(z) A^T(q)^{-1}, \quad (6)$$

where:

$$A^T(q) = \begin{bmatrix} q^{\frac{1}{8}} & 0 \\ 0 & q^{-\frac{1}{8}} \end{bmatrix}. \quad (7)$$

The gauge transformation acts in the "internal space" of the L-operator too, i.e. in addition to conjugation (6) we should change the dynamical variables:

$$\begin{bmatrix} S_3 & 2S^+ \\ 2S^- & -S_3 \end{bmatrix} \rightarrow \begin{bmatrix} q^{\frac{1}{8}} & 0 \\ 0 & q^{-\frac{1}{8}} \end{bmatrix}^{-1} \begin{bmatrix} S_3 & 2S^+ \\ 2S^- & -S_3 \end{bmatrix} \begin{bmatrix} q^{\frac{1}{8}} & 0 \\ 0 & q^{-\frac{1}{8}} \end{bmatrix},$$

Or more explicitly:

$$S_3 \rightarrow S_3, \quad S^+ \rightarrow q^{-\frac{1}{4}} S^+, \quad S^- \rightarrow q^{\frac{1}{4}} S^-.$$

After applying the gauge transformation of such a form, we can take the limit $\tau \rightarrow i\infty$ in (6). We get:

$$\left\{ \begin{array}{l} S_3 = -\frac{v}{u \tan(2\pi u)} - \frac{\nu}{\sin^2(2\pi u)}, \\ S^+ = -\frac{v}{4\pi \sin(2\pi u)} - \frac{\nu \cos(2\pi u)}{4 \sin^2(2\pi u)}, \\ S^- = \frac{v \cos^2(2\pi u)}{\pi \sin(2\pi u)} + \frac{\nu \cos(2\pi u) (1 + \sin^2(2\pi u))}{\sin^2(2\pi u)}. \end{array} \right. \quad (8)$$

With respect to the canonical Poisson bracket $\{u, v\} = 1$ we have:

$$\{S_3, S^+\} = 2S^+, \quad \{S_3, S^-\} = -2S^-, \quad \{S^+, S^-\} = S_3.$$

The Casimir function:

$$\Omega = S_3^2 + 4S^+ S^- = \nu^2.$$

Equations (8) can be presented in a form:

$$L^{TT}(z) = \Xi^T(z) L^{TKM}(z) \Xi^T(z)^{-1}, \quad (9)$$

where:

$$L^{TT}(z) = \lim_{q \rightarrow 0} A^T(q) L^{ET}(z) A^T(q)^{-1}, \quad \Xi^T(z) = q^{\frac{1}{8}} \lim_{q \rightarrow 0} A^T(q) \Xi^E(z), \quad L^{TKM}(z) = \lim_{q \rightarrow 0} L^{ECM}(z).$$

Taking the limits we get:

$$\begin{aligned} L^{TKM}(z) &= \begin{bmatrix} v & \nu \pi (\cot(2\pi u) + \cot(\pi z)) \\ \nu \pi (-\cot(2\pi u) + \cot(\pi z)) & -v \end{bmatrix}, \\ \Xi^T(z) &= \begin{bmatrix} 1 & -1 \\ -2 \cos(\pi(z - 2u)) & 2 \cos(\pi(z + 2u)) \end{bmatrix}, \\ L^{TT}(z) &= \begin{bmatrix} \pi \cot(\pi z) S_3 & \frac{2\pi S^+}{\sin(\pi z)} \\ \frac{2\pi S^-}{\sin(\pi z)} + 8\pi S^+ \sin(\pi z) & -\pi \cot(\pi z) S_3 \end{bmatrix}. \end{aligned} \quad (10)$$

It is worth noting that L-operator (10) does not coincide with the standard trigonometric degeneration of the elliptical one. Thus, this Lax matrix corresponds to some new trigonometric solution of the Yang-Baxter equation. Let us calculate the hamiltonian for the system defined by $L^{TT}(z)$:

$$\text{Tr}(L^{TT}(z)^2) = 2\pi^2 (S_3^2 - 16S^{+2}) - \frac{2\pi^2 \Omega}{\sin^2(\pi z)},$$

and from the definition for the hamiltonian we have:

$$H^T = \frac{1}{4\pi i} \oint \frac{\text{tr} L^{TT}(z)^2}{z} dz = \pi^2 (S_3^2 - 16S^{+2}). \quad (11)$$

Using (8) we see that this hamiltonian describes a motion of the trigonometric CM system:

$$H^T = \pi^2 (S_3^2 - 16 S^{+2}) = v^2 - \frac{\pi^2 \nu^2}{\sin^2(2\pi u)}.$$

Let us move to the rational case. To get the rational degenerations of (8) we should replace the constant π (the half period of the trigonometric functions) by a new variable x and then take the limit as x goes to zero. Just as in the trigonometric case we see that in this limit the right hand side of (8) diverges. To avoid this difficulty we apply to (9) an additional gauge transformation:

$$A^R(x) L^{TT}(z) A^R(x)^{-1} = A^R(x) \Xi^T(z) L^{TCM}(z) \Xi^T(z)^{-1} A^R(x)^{-1}, \quad (12)$$

where the matrix $A^R(x)$ has the form:

$$A^R(x) = \begin{bmatrix} x & 0 \\ \frac{2}{x} & \frac{1}{x} \end{bmatrix}. \quad (13)$$

The gauge transformation in the internal space acts as follows:

$$\begin{bmatrix} S_3 & 2S^+ \\ 2S^- & -S_3 \end{bmatrix} \longrightarrow \begin{bmatrix} x & 0 \\ \frac{2}{x} & \frac{1}{x} \end{bmatrix}^{-1} \begin{bmatrix} S_3 & 2S^+ \\ 2S^- & -S_3 \end{bmatrix} \begin{bmatrix} x & 0 \\ \frac{2}{x} & \frac{1}{x} \end{bmatrix},$$

i.e. we have a substitution:

$$S_3 \rightarrow S_3 + \frac{4S^+}{x^2}, \quad S^+ \rightarrow \frac{S^+}{x^2}, \quad S^- \rightarrow S^- x^2 - 2S_3 - \frac{4S^+}{x^2}.$$

After such a transformation we can take the limit $x = 0$ in (12) and we find:

$$\begin{cases} S_3 = uv - \frac{\nu}{2}, \\ S^+ = \frac{v}{2u} + \frac{\nu}{4u^2}, \\ S^- = -\frac{u^3 v}{2} + \frac{3\nu u^2}{4}. \end{cases} \quad (14)$$

With respect to the canonical Poisson bracket $\{u, v\} = 1$ we have:

$$\{S_3, S^+\} = 2S^+, \quad \{S_3, S^-\} = -2S^-, \quad \{S^+, S^-\} = S_3, \quad \Omega = S_3^2 + 4S^+ S^- = \nu^2$$

Expressions (14) can be presented in a form similar to (3):

$$L^{RT}(z) = \Xi^R(z) L^{RCM}(z) \Xi^R(z)^{-1}, \quad (15)$$

where

$$\begin{aligned} L^{RCM}(z) &= \begin{bmatrix} v & \nu \left(\frac{1}{z} + \frac{1}{2u} \right) \\ \nu \left(\frac{1}{z} - \frac{1}{2u} \right) & -v \end{bmatrix}, \\ \Xi^R(z) &= \begin{bmatrix} -1 & 1 \\ -zu + u^2 & -zu - u^2 \end{bmatrix}, \\ L^{RT}(z) &= \begin{bmatrix} \frac{S_3}{z} & \frac{2S^+}{z} \\ \frac{2S^-}{z} + zS_3 & -\frac{S_3}{z} \end{bmatrix}. \end{aligned} \quad (16)$$

Lax matrix (16) is a new rational L-operator for $\mathfrak{sl}(2, \mathbb{C})$ because it does not coincide with the standard rational one. This Lax matrix defines some new hamiltonian system on the algebra $\mathfrak{sl}(2, \mathbb{C})$. The hamiltonian of this system can be obtained from the expansion:

$$\text{tr} L^{RT}(z)^2 = -4S^+ S_3 + \frac{2\Omega}{z^2} \Rightarrow H^R = 2S^+ S_3.$$

Using (14) we see that this hamiltonian coincides with the hamiltonian of the rational CM system:

$$H^R = 2 S^+ S_3 = v^2 - \frac{\nu^2}{4u^2}. \quad (17)$$

Thus, we have shown that there exist new trigonometrical and rational L-operators corresponding to the two particle CM systems. These new Lax operators can be obtained from the elliptical one by applying to it some additional gauge transformations. One can generalize this construction to an arbitrary rank. A main difficulty here is to find the matrices of the additional gauge transformations $A^T(q)$ and $A^R(x)$ for all N .

3 The Elliptic $SL(N)$ Top

The elliptic $sl(N, \mathbb{C})$ top is the integrable system defined by the classical Belavin-Drinfeld L-operator:

$$L^{ET} = \sum_{n+m \neq 0}^{N-1} S_{mn} \varphi_{m,n}(z) T_{mn}. \quad (18)$$

Here $\varphi_{m,n}(z)$ are defined by the standard theta functions (2):

$$\varphi_{m,n}(z) = \exp\left(-\frac{2\pi i n z}{N}\right) \phi\left(-\frac{m+n\tau}{N}; z\right), \quad \phi(u, z) = \frac{\theta_{11}(u+z) \theta'_{11}(0)}{\theta_{11}(u) \theta_{11}(z)}. \quad (19)$$

where S_{mn} are the generalized coordinates of the top and T_{mn} are the $N \times N$ matrices of the $sl(N, \mathbb{C})$ generators in the fundamental representation. The generators T_{mn} correspond to the so called sin-alg basis and can be constructed as follows:

$$T_{mn} = \exp\left(\frac{\pi i m n}{N}\right) Q^m \Lambda^n, \quad (m = 0 \dots N-1, n = 0 \dots N-1, n+m \neq 0),$$

$$Q = \text{diag} \left[\exp\left(\frac{2\pi i}{N}\right), \exp\left(\frac{4\pi i}{N}\right), \dots, \exp\left(\frac{2m\pi i}{N}\right), \dots, 1 \right],$$

$$\Lambda = \begin{bmatrix} 0 & 1 & 0 & \dots & \dots & 0 \\ 0 & 0 & 1 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \dots & 1 \\ 1 & 0 & 0 & \dots & \dots & 0 \end{bmatrix}.$$

It is easy to see that these operators form the $sl(N, \mathbb{C})$ basis, and obey the following relations:

$$[T_{sk}, T_{nj}] = 2i \sin\left(\frac{\pi}{N}(kn - sj)\right) T_{[s+n], [k+j]}, \quad Tr(T_{sk} T_{nj}) = \delta_{s, -n} \delta_{k, -j} N. \quad (20)$$

Here $|a|$ is the value of a modulo N that belongs to the interval $(1..N)$, i.e. $|0| = N$. The dynamics of the top is defined by the Lax equation by means of the auxiliary M-matrix:

$$\partial_t L^E(z) = [M^E(z), L^E(z)], \quad M^E(z) = \sum_{n+m \neq 0}^{N-1} \exp\left(-\frac{nz}{N}\right) f\left(\frac{-m+n\tau}{N}, z\right) S_{mn} T_{mn}, \quad f(u, z) = \partial_u \phi(u, z).$$

From Lax equation we can get the equations of motion:

$$\partial_t S_{mn} = \frac{N}{\pi} \sum_{kl} S_{kl} S_{m-k, n-l} \wp\left(\frac{k+l\tau}{N}, \tau\right) \sin \frac{\pi}{N}(kn - ml), \quad (21)$$

where $\wp(z, \tau)$ is the Weierstrass elliptic function. The quadratic hamiltonian of the system is defined by the expansion:

$$\frac{1}{N} \text{tr} L^E(z)^2 = -2 H^E + \Omega_2 \wp(z, \tau), \quad \Omega_2 = \sum_{n+m \neq 0}^{N-1} S_{mn}^2, \quad (22)$$

where

$$H^E = -\frac{1}{2} \sum_{m+n \neq 0}^{N-1} \wp\left(\frac{m+n\tau}{N}, \tau\right) S_m S_{-m-n}. \quad (23)$$

In [5] the correspondence between the elliptic $\mathfrak{sl}(N, \mathbb{C})$ top and the N -particle elliptic CM system was established. The Lax operators of these systems are connected by the equation:

$$L^{ET} = \Xi(z) L^{ECM}(z) \Xi(z)^{-1}, \quad (24)$$

where

$$\begin{aligned} L^{ECM}(z) &= v_i \delta_{ij} + (1 - \delta_{ij}) \nu \phi(u_j - u_i, z), \\ \Xi(z) &= \Xi'(z) \times \text{diag} \left((-1)^l \left(\prod_{j < k, j, k \neq l} \theta_{11}(u_k - u_j, \tau) \right)^{-1} \right), \\ \Xi'(z)_{ij} &= \theta_{\frac{i}{N} - \frac{1}{2}, \frac{N}{2}}(z - Nu_j, N\tau). \end{aligned}$$

4 The Trigonometric Lax Operator

The trigonometric degeneration corresponds to the limit $q \rightarrow 0$. It can be checked that in this limit the right-hand side of (24) diverges and as well as in the $\mathfrak{sl}(2, \mathbb{C})$ case we need to apply to this equation some additional regularising gauge transformation depending on q . To find the matrix $A^T(q)$ of the additional gauge transformation let us consider behaviour of the matrix elements of $\Xi(z)$ in the neighbourhood of the point $q = 0$. We find that:

$$\Xi_{i,j} \sim q^{\frac{N}{2} \left(\frac{i}{N} - \frac{1}{2} \right)^2}, \quad \Xi_{i,j}^{-1} \sim q^{-\frac{N}{2} \left(\frac{j}{N} - \frac{1}{2} \right)^2}, \quad q = \exp(2\pi i \tau). \quad (25)$$

From these expressions we find the leading terms for the right hand side of (24):

$$R_{ij} = \Xi_{i,k}(z) L_{k,s}^{CM}(z) \Xi_{sj}^{-1}(z) \sim q^{\frac{N}{2} \left(\left(\frac{i^2}{N^2} - \frac{i}{N} \right) - \left(\frac{j^2}{N^2} - \frac{j}{N} \right) \right)}. \quad (26)$$

Analogously to the $\mathfrak{sl}(2, \mathbb{C})$ case let us consider the matrix $A^T(q)$ in the diagonal form:

$$A^T(q) = \text{diag} [q^{a_1}, q^{a_2} \dots q^{a_N}]. \quad (27)$$

The degrees a_i are defined from the condition for regularity of (24) after applying the the gauge transformation $A^T(q)$ in the limit $q = 0$. Combining (26) and (27) we have:

$$[A^T(q) R A^T(q)^{-1}]_{ij} \sim q^{a_i - a_j + \frac{N}{2} \left(\left(\frac{i^2}{N^2} - \frac{i}{N} \right) - \left(\frac{j^2}{N^2} - \frac{j}{N} \right) \right)}.$$

Therefore, the necessary condition for a_i has the following form:

$$a_j - a_i = \frac{N}{2} \left(\left(\frac{i^2}{N^2} - \frac{i}{N} \right) - \left(\frac{j^2}{N^2} - \frac{j}{N} \right) \right). \quad (28)$$

The second condition is that the matrix $A^T(q)$ belongs to the gauge group $\text{SL}(N)$ for all q :

$$\det A^T(q) = 1, \quad \Rightarrow \quad \sum_{i=1}^N a_i = 0. \quad (29)$$

From (28) and (29) we get:

$$a_i = - \left(\frac{N}{2} \left(\frac{i^2}{N^2} - \frac{i}{N} \right) + \frac{N^2 - 1}{12N} \right). \quad (30)$$

After constructing the matrix $A^T(q)$ we can find the trigonometric Lax operator:

$$L^{TT}(z) = \lim_{q \rightarrow 0} A(q) L^{ET}(z) A^{-1}(q) \quad (31)$$

Here as well as in the $\mathfrak{sl}(2, \mathbb{C})$ case in addition to conjugation (31) the gauge transformation acts in the internal space of the L-operator. To alleviate the calculations it is useful to work in the basis g_{ij} induced by the natural embedding $\mathfrak{sl}(N, \mathbb{C}) \hookrightarrow \mathfrak{gl}(N, \mathbb{C})$. The connection between the sin-alg basis and $\mathfrak{gl}(N, \mathbb{C})$ basis has the form:

$$S_{n,m} = \sum_{|b-a|=n} \exp\left(-\frac{\pi i m(|b-a| + 2|a|)}{N}\right) g_{ab}, \quad \{g_{ij}, g_{km}\} = \delta_{jk} g_{im} - \delta_{im} g_{kj}.$$

To give the answer for the limit (31), which seems complicated, it is useful to write it down by parts. For the nondiagonal part we get:

$$\left\{ \begin{array}{ll} a < b & \begin{aligned} L_{a,b}^{TT} &= \frac{N\pi}{\sin(\pi z)} \exp\left(-\frac{\pi i z}{N}(N-2b+2a)\right) g_{a,b-} \\ &+ 2\pi i N \exp\left(\frac{2\pi i(b-a)z}{N}\right) \sum_{k=1}^{N-b} g_{a+k,b+k}, \end{aligned} \\ a > b, a \neq N & \begin{aligned} L_{a,b}^{TT} &= \frac{N\pi}{\sin(\pi z)} \exp\left(-\frac{\pi i z}{N}(-N-2b+2a)\right) g_{ab+} \\ &+ 2\pi N i \exp\left(-\frac{2\pi i(a-b)z}{N}\right) \sum_{k=1}^b g_{a-k,|b-k|}, \end{aligned} \\ a = N & \begin{aligned} L_{N,b}^{TT} &= \frac{N\pi}{\sin(\pi z)} \exp\left(-\frac{\pi i(N-2b)z}{N}\right) g_{Nb-} \\ &+ 2\pi i N \exp\left(\frac{2\pi i b z}{N}\right) \sum_{k=1}^{N-b} g_{k,b+k} + \\ &+ 2\pi i N \exp\left(\frac{2\pi i(b-N)z}{N}\right) \sum_{k=0}^{b-1} g_{|N-b+k|,|N+k|}. \end{aligned} \end{array} \right. , \quad (32)$$

The diagonal part of $L^{TT}(z)$ is:

$$L_{a,a}^{TT} = \frac{\pi}{2 \sin(\pi z)} [A_a \exp(-\pi i z) + B_a \exp(\pi i z)], \quad (33)$$

where:

$$A_a = \sum_{k=1}^N (N-1-2|k-a|) g_{k,k}, \quad B_a = \sum_{k=1}^N (N-1-2|a-k|) g_{k,k}. \quad (34)$$

The connection with the N-particle trigonometric CM system has the form:

$$L^{TT}(z) = \Xi^T(z) L^{TCM}(z) \Xi^T(z)^{-1}. \quad (35)$$

where $L^{TCM}(z)$ is the Lax operator for the N-particle trigonometric CM system:

$$L^{TCM}(u)_{ij} = v_i \delta_{ij} + (1 - \delta_{ij}) \nu \pi (\cot(\pi z) + \cot(\pi(u_i - u_j))) \quad (36)$$

and the matrix $\Xi^T(z)$ is defined as the limit:

$$\Xi^T(z) = \lim_{q \rightarrow 0} q^{\frac{N^2-1}{12N}} A^T(q) \Xi(z). \quad (37)$$

Lax matrix (32) defines some new integrable system on the algebra $\mathfrak{sl}(N, \mathbb{C})$. The hamiltonian for this system is defined in the standard way:

$$H^T = \frac{1}{2\pi i} \oint \frac{\text{tr} L^{TT}(z)^2}{z} dz.$$

Using equation (35) for the hamiltonian we get:

$$H^T = \frac{1}{2\pi i} \oint \frac{\text{tr}(L^{TT}(z)^2)}{z} dz = \frac{1}{2\pi i} \oint \frac{\text{tr}(L^{TCM}(z)^2)}{z} dz = \sum_{n=1}^N v_n^2 + \sum_{i \neq j}^N \frac{\pi^2 \nu^2}{\sin^2 \pi(u_i - u_j)} = H^{TCM}. \quad (38)$$

Therefore, we get the equivalence between the N-particle trigonometric CM system and some new integrable top on the algebra $\mathfrak{sl}(N, \mathbb{C})$ defined by L-operator (32). It worth noting that this Lax has very sophisticated form, and it does not coincide with the standard trigonometric L-operator for the algebra $\mathfrak{sl}(N, \mathbb{C})$. To illustrate the formula (32) we give in appendix the explicit expressions for $L^{TT}(z)$ in the cases $N=2,3,4$. We calculate the integrals of motion for the system defined by these Lax and check that they commute.

5 The Rational Case

The rational degeneration corresponds to the limit as the period of the trigonometric functions approaches infinity. To find this degeneration we replace in (35) the constant π by a new variable x and then tend x to zero. As well as in the $\text{sl}(2, \mathbb{C})$ case the right-hand side of (35) diverges in this limit. Analogously to the trigonometric case we apply to this equation the regularizing gauge transformation depending on the parameter x . Let us consider the matrix $A^R(x)$ for this transformation in the form:

$$A^R(x) = W_2 W_1, \quad (39)$$

here the matrix W_1 is lower-triangular and does not depend on x . The matrix W_2 is diagonal:

$$W_2 = \text{diag}(x^{b_1}, x^{b_2}, \dots, x^{b_N}). \quad (40)$$

As in the previous section we need to find the degrees b_i and the coefficients of W_1 from the condition for regularity of the limit:

$$\lim_{x \rightarrow 0} A^R(x) \Xi^T(u) L^{TCM}(z) \Xi^T(u)^{-1} A^R(x)^{-1}.$$

We find:

$$b_i = \left(-\frac{N(N-1)}{2N} + \frac{1-(i-1)N}{N} \right) \quad (41)$$

$$\begin{cases} W_{1ij} = \frac{(i-1)!}{(j-i)!(j-1)!}, & i \neq N, j \leq i, \\ W_{1Nj} = \frac{N!}{i!(N-i)!}, & j \in (1 \dots N) \\ W_{1ij} = 0, & j > i \end{cases} \quad (42)$$

Having found the regularizing matrix $A^R(x)$ we can get the rational version of (35):

$$L^{RT}(z) = \Xi^R(u) L^{RCM}(z) \Xi^R(u)^{-1}. \quad (43)$$

where $L^{RCM}(z)$ is the Lax operator for the rational N-particle CM system:

$$L^{RCM}(u)_{ij} = v_i \delta_{ij} + \nu (1 - \delta_{ij}) \frac{\nu}{u_i - u_j},$$

and the matrices $\Xi^R(z)$ and $L^{RT}(z)$ are defined as the limits:

$$\Xi^R(z) = \lim_{x \rightarrow 0} x^{K_N} A^R(x) \Xi(z), \quad L^{RT}(z) = \lim_{x \rightarrow 0} A^R(x) L^{TT}(z) A^R(x)^{-1}, \quad (44)$$

where K_N is some constant defined from the condition for regularity of the limit for $\Xi^R(z)$. The hamiltonian of the rational top is defined in the standard way:

$$H^R = \frac{1}{2\pi i} \oint \frac{\text{tr} L^{RT}(z)^2}{z} dz.$$

Using (43) for this hamiltonian we get:

$$H^R = \frac{1}{2\pi i} \oint \frac{\text{tr} L^{RT}(z)^2}{z} dz = \frac{1}{2\pi i} \oint \frac{\text{tr} L^{RCM}(z)^2}{z} dz = \sum_{n=1}^N v_n^2 + \sum_{i \neq j}^N \frac{\nu^2}{(u_i - u_j)^2} = H^{RCM}. \quad (45)$$

The last equation gives the equivalence between the rational top on the algebra $\text{sl}(N, \mathbb{C})$ defined by Lax operator (43) and the N-particle rational CM system. In this section we have not presented the explicit expression for the $L^{RT}(z)$. Nevertheless, using the matrix of additional gauge transformation (39) by means of modern computational tools such as Maple or Mathematica we can calculate limits (44) for $N \leq 10$. In appendix we present the examples of such a calculations for $N=2,3,4$. We give the explicit expressions for the L-operators corresponding to these cases and calculate the integrals of motion for the rational tops and then check that they commute.

Acknowledgments

The author is grateful to M. Olshanetsky, A. Levin, A. Zotov and A. Sleptsov for fruitful discussions and interest to this work. The work was partly supported by RFBR grant 06-02-17382, RFBR grant 06-01-92054- KE_a and grant for support of scientific schools NSh-3036.2008.2.

Appendix T

Here we give the examples of the trigonometrical Lax operators for $N=2,3,4$. We use the following normalization for Lax operators:

$$L^{TT}(z) = \frac{\pi N}{\sin(\pi z)} L(z).$$

The operator $L(z)$ for $N = 2$:

$$L = -1/2 \cos(\pi z) (-g_{1,1} + g_{2,2})$$

$$L_{12} = g_{1,2}$$

$$L_{21} = -4 g_{1,2} (\cos(\pi z))^2 + 4 g_{1,2} + g_{2,1}$$

$$L_{22} = 1/2 \cos(\pi z) (-g_{1,1} + g_{2,2})$$

The operator $L(z)$ for $N = 3$:

$$L_{11} = (-e^{-i\pi z} g_{3,3} + e^{-i\pi z} g_{1,1} - e^{i\pi z} g_{2,2} + e^{i\pi z} g_{1,1})/3$$

$$L_{12} = e^{-1/3 i\pi z} g_{1,2} - g_{2,3} e^{5/3 i\pi z} + e^{-1/3 i\pi z} g_{2,3}$$

$$L_{13} = g_{1,3} e^{1/3 i\pi z}$$

$$L_{21} = g_{2,1} e^{1/3 i\pi z} + g_{1,3} e^{1/3 i\pi z} - e^{-5/3 i\pi z} g_{1,3}$$

$$L_{22} = (e^{-i\pi z} g_{2,2} - e^{-i\pi z} g_{1,1} - e^{i\pi z} g_{3,3} + e^{i\pi z} g_{2,2})/3$$

$$L_{23} = e^{-1/3 i\pi z} g_{2,3}$$

$$L_{31} = -g_{2,3} e^{5/3 i\pi z} + e^{-1/3 i\pi z} g_{1,2} - g_{1,2} e^{5/3 i\pi z} - e^{-7/3 i\pi z} g_{2,3} + g_{3,1} e^{-1/3 i\pi z} + 2 e^{-1/3 i\pi z} g_{2,3}$$

$$L_{32} = g_{3,2} e^{1/3 i\pi z} - g_{1,3} e^{7/3 i\pi z} + 2 g_{1,3} e^{1/3 i\pi z} - e^{-5/3 i\pi z} g_{2,1} - e^{-5/3 i\pi z} g_{1,3} + g_{2,1} e^{1/3 i\pi z}$$

$$L_{33} = (-e^{i\pi z} g_{1,1} + e^{-i\pi z} g_{3,3} - e^{-i\pi z} g_{2,2} + e^{i\pi z} g_{3,3})/3$$

The operator $L(z)$ for $N = 4$:

$$L_{11} = (-e^{i\pi z} g_{3,3} + 3 e^{-i\pi z} g_{1,1} - 3 e^{-i\pi z} g_{4,4} + e^{-i\pi z} g_{2,2} - 3 e^{i\pi z} g_{2,2} + 3 e^{i\pi z} g_{1,1} + e^{i\pi z} g_{4,4} - e^{-i\pi z} g_{3,3})/8$$

$$L_{12} = e^{-1/2 i\pi z} g_{3,4} + e^{-1/2 i\pi z} g_{1,2} - g_{3,4} e^{3/2 i\pi z} + e^{-1/2 i\pi z} g_{2,3} - g_{2,3} e^{3/2 i\pi z}$$

$$L_{13} = g_{2,4} + g_{1,3} - g_{2,4} e^{2 i\pi z}$$

$$L_{14} = g_{1,4} e^{1/2 i\pi z}$$

$$L_{21} = g_{1,4} e^{1/2 i\pi z} - e^{-3/2 i\pi z} g_{1,4} + g_{2,1} e^{1/2 i\pi z}$$

$$L_{22} = (-3 e^{i\pi z} g_{3,3} - 3 e^{-i\pi z} g_{1,1} - e^{-i\pi z} g_{4,4} + 3 e^{-i\pi z} g_{2,2} + 3 e^{i\pi z} g_{2,2} + e^{i\pi z} g_{1,1} - e^{i\pi z} g_{4,4} + e^{-i\pi z} g_{3,3})/8$$

$$L_{23} = -g_{3,4}e^{3/2 i\pi z} + e^{-1/2 i\pi z}g_{2,3} + e^{-1/2 i\pi z}g_{3,4}$$

$$L_{24} = g_{2,4}$$

$$L_{31} = g_{2,4} - e^{-2 i\pi z}g_{2,4} + g_{3,1}$$

$$L_{32} = -e^{-3/2 i\pi z}g_{2,1} + g_{1,4}e^{1/2 i\pi z} - e^{-3/2 i\pi z}g_{1,4} + g_{2,1}e^{1/2 i\pi z} + g_{3,2}e^{1/2 i\pi z}$$

$$L_{33} = (3e^{i\pi z}g_{3,3} - e^{-i\pi z}g_{1,1} + e^{-i\pi z}g_{4,4} - 3e^{-i\pi z}g_{2,2} + e^{i\pi z}g_{2,2} - e^{i\pi z}g_{1,1} - 3e^{i\pi z}g_{4,4} + 3e^{-i\pi z}g_{3,3})/8$$

$$L_{34} = e^{-1/2 i\pi z}g_{3,4}$$

$$L_{41} = -g_{2,3}e^{3/2 i\pi z} + 2e^{-1/2 i\pi z}g_{3,4} + e^{-1/2 i\pi z}g_{2,3} + g_{4,1}e^{-1/2 i\pi z} - e^{-5/2 i\pi z}g_{3,4} - g_{1,2}e^{3/2 i\pi z} - g_{3,4}e^{3/2 i\pi z} + e^{-1/2 i\pi z}g_{1,2}$$

$$L_{42} = -e^{-2 i\pi z}g_{2,4} + 2g_{2,4} + g_{3,1} + g_{1,3} - g_{2,4}e^{2 i\pi z} - g_{1,3}e^{2 i\pi z} + g_{4,2} - e^{-2 i\pi z}g_{3,1}$$

$$L_{43} = 2g_{1,4}e^{1/2 i\pi z} - e^{-3/2 i\pi z}g_{1,4} - g_{1,4}e^{5/2 i\pi z} - e^{-3/2 i\pi z}g_{2,1} + g_{4,3}e^{1/2 i\pi z} + g_{2,1}e^{1/2 i\pi z} - e^{-3/2 i\pi z}g_{3,2} + g_{3,2}e^{1/2 i\pi z}$$

$$L_{44} = (e^{i\pi z}g_{3,3} + e^{-i\pi z}g_{1,1} + 3e^{-i\pi z}g_{4,4} - e^{-i\pi z}g_{2,2} - e^{i\pi z}g_{2,2} - 3e^{i\pi z}g_{1,1} + 3e^{i\pi z}g_{4,4} - 3e^{-i\pi z}g_{3,3})/8$$

The hamiltonians and highest integrals of motion

On calculating the integrals of motion we use the standard definition:

$$H_k = \frac{1}{2k\pi i} \oint \frac{\text{tr}(L^{TT}(z))^k}{z} dz.$$

The integral H_2 is usually identified with the hamiltonian. We check the commutativity with respect to the standard Poisson bracket on the algebra $\text{gl}(N, \mathbb{C})$:

$$\{A, B\}_N = \sum_{i,j,k=1}^N \left(\frac{\partial A}{\partial g_{i,j}} \frac{\partial B}{\partial g_{j,k}} - \frac{\partial B}{\partial g_{i,j}} \frac{\partial A}{\partial g_{j,k}} \right) g_{i,k}$$

The case N=2

$$H_2 = 1/2 (g_{2,2} - g_{1,1} + 4g_{1,2})(g_{2,2} - g_{1,1} - 4g_{1,2})\pi^2$$

The case N=3

$$H_2 = 2 (g_{2,2}^2 - g_{2,2}g_{3,3} - 9g_{2,3}g_{2,1} - g_{1,1}g_{3,3} - 9g_{1,2}g_{1,3} + g_{1,1}^2 - 27g_{1,3}g_{2,3} - g_{1,1}g_{2,2} + g_{3,3}^2)\pi^2$$

$$H_3 = g_{1,1}^2g_{2,2} + 27g_{1,3}^2g_{2,1} - g_{1,1}^2g_{3,3} + g_{2,2}^2g_{3,3} - 9g_{1,2}g_{2,2}g_{1,3} - 27g_{2,3}^3 + 9g_{2,3}g_{2,1}g_{1,1} - g_{1,1}g_{2,2}^2 + g_{1,1}g_{3,3}^2 + 27g_{1,3}^3 - 27g_{2,3}^2g_{1,2} - 9g_{3,3}g_{2,3}g_{2,1} + 9g_{1,2}g_{1,3}g_{3,3} - g_{2,2}g_{3,3}^2$$

And we have checked that

$$\{H_2, H_2\}_3 = 0.$$

The case N=4

$$H_2 = 2g_{1,1}g_{2,2} + 6g_{1,1}g_{3,3} - 5g_{4,4}^2 + 6g_{4,4}g_{2,2} + 2g_{4,4}g_{3,3} + 32g_{3,4}g_{3,2} + 2g_{2,2}g_{3,3} + 32g_{1,3}g_{2,4} + 32g_{2,3}g_{2,1} + 64g_{2,4}^2 + 32g_{2,4}g_{3,1} - 5g_{1,1}^2 + 2g_{1,1}g_{4,4} + 64g_{3,4}g_{2,1} + 128g_{1,4}g_{3,4} - 5g_{2,2}^2 + 32g_{1,2}g_{1,4} - 5g_{3,3}^2 + 64g_{2,3}g_{1,4}$$

$$H_3 = 4g_{1,3}g_{2,4}g_{4,4} - 4g_{1,3}g_{2,4}g_{3,3} + 4g_{1,3}g_{2,4}g_{2,2} - 16g_{3,4}^2g_{1,3} - g_{1,1}^2g_{4,4} - 4g_{1,3}g_{2,4}g_{1,1} - 64g_{3,4}g_{2,3}g_{2,4} - 32g_{1,2}g_{3,4}g_{2,4} - 8g_{3,4}g_{2,1}g_{4,4} - 16g_{3,4}g_{2,3}g_{3,1} + 4g_{2,4}g_{3,1}g_{1,1} + 8g_{3,4}g_{2,1}g_{1,1} - 8g_{2,3}g_{2,1}g_{3,3} + 4g_{2,4}g_{3,1}g_{3,3} - 4g_{2,4}g_{3,1}g_{2,2} - 4g_{2,4}g_{3,1}g_{4,4} - 8g_{3,4}g_{3,2}g_{4,4} + 8g_{1,2}g_{1,4}g_{4,4} - 8g_{3,4}g_{2,1}g_{3,3} - 8g_{2,3}g_{1,4}g_{3,3} + 8g_{2,3}g_{2,1}g_{1,1} - 16g_{1,2}g_{2,3}g_{2,4} + 8g_{2,3}g_{1,4}g_{1,1} + 8g_{3,4}g_{3,2}g_{2,2} + 8g_{2,3}g_{1,4}g_{4,4} - 8g_{1,2}g_{1,4}g_{2,2} + 16g_{1,3}g_{2,1}g_{1,4} + 8g_{3,4}g_{2,1}g_{2,2} + 64g_{2,4}g_{2,1}g_{1,4} + 32g_{2,4}g_{3,2}g_{1,4} - 8g_{2,3}g_{1,4}g_{2,2} + 16g_{1,3}g_{1,4}^2 + 16g_{2,4}g_{2,1}^2 + 64g_{2,4}g_{1,4}^2 - 16g_{3,4}^2g_{3,1} - 16g_{2,3}^2g_{2,4} + 16g_{1,4}^2g_{3,1} - 64g_{3,4}^2g_{2,4} + g_{2,2}^2g_{3,3} - g_{2,2}g_{3,3}^2 + g_{4,4}g_{3,3}^2 - g_{1,1}g_{2,2}^2 +$$

$$g_{1,1}g_{4,4}^2 - g_{4,4}^2g_{3,3} + 16g_{2,4}g_{3,2}g_{2,1} + g_{1,1}^2g_{2,2}$$

$$\begin{aligned} H_4 = & 49152g_{2,4}^2g_{1,4}g_{3,4} + 3584g_{4,4}g_{2,2}g_{3,4}g_{1,4} + 1536g_{4,4}g_{1,2}g_{2,3}g_{2,4} - 1536g_{2,4}g_{3,2}g_{2,1}g_{1,1} + 8192g_{1,3}g_{2,4}g_{2,3}g_{1,4} \\ & - 256g_{3,4}g_{2,1}g_{1,1}g_{4,4} + 512g_{2,2}g_{3,3}g_{1,4}g_{3,4} + 4096g_{1,3}g_{2,1}g_{2,3}g_{2,4} - 2048g_{2,4}g_{2,2}g_{3,2}g_{1,4} + 4096g_{2,4}g_{3,2}g_{3,4}g_{3,1} + 1792g_{4,4}g_{2,2}g_{2,3}g_{1,4} + \\ & 1536g_{3,3}g_{1,3}g_{2,1}g_{1,4} - 128g_{1,1}g_{2,2}g_{3,4}g_{3,2} + 384g_{4,4}g_{3,3}g_{3,4}g_{3,2} + 4096g_{2,3}g_{2,1}g_{2,4}g_{3,1} + 384g_{2,3}g_{2,1}g_{2,2}g_{3,3} + 3584g_{1,1}g_{3,3}g_{3,4}g_{1,4} + \\ & 384g_{2,2}g_{3,3}g_{3,4}g_{3,2} + 512g_{1,3}g_{2,1}g_{1,4}g_{2,2} - 128g_{2,3}g_{2,1}g_{4,4}g_{3,3} - 512g_{2,2}g_{1,2}g_{2,3}g_{2,4} + 384g_{1,1}g_{3,3}g_{1,2}g_{1,4} - 1536g_{3,4}g_{2,3}g_{3,1}g_{4,4} - \\ & 128g_{4,4}g_{3,3}g_{1,2}g_{1,4} + 384g_{1,1}g_{2,2}g_{1,2}g_{1,4} + 24576g_{3,4}g_{2,3}g_{2,1}g_{1,4} + 1408g_{4,4}g_{2,2}g_{3,4}g_{3,2} + 256g_{3,4}g_{2,1}g_{4,4}g_{3,3} + 4096g_{3,4}g_{2,3}g_{3,2}g_{1,4} - \\ & 512g_{2,4}g_{2,2}g_{3,2}g_{2,1} + 1408g_{4,4}g_{2,2}g_{1,2}g_{1,4} + 768g_{3,4}g_{2,1}g_{2,2}g_{3,3} + 4096g_{3,4}g_{1,3}g_{3,2}g_{2,4} + 4096g_{3,4}g_{2,3}g_{3,2}g_{2,1} + 256g_{1,1}g_{4,4}g_{2,3}g_{1,4} + \\ & 1536g_{2,4}g_{4,4}g_{3,2}g_{2,1} + 384g_{1,1}g_{3,3}g_{3,4}g_{3,2} + 2048g_{4,4}g_{3,4}g_{2,3}g_{2,4} + 1408g_{2,3}g_{2,1}g_{1,1}g_{3,3} - 128g_{2,3}g_{2,1}g_{1,1}g_{4,4} + 384g_{2,3}g_{2,1}g_{4,4}g_{2,2} + \\ & 4096g_{1,3}g_{2,4}g_{1,2}g_{1,4} + 1536g_{1,1}g_{2,3}g_{3,4}g_{3,1} - 2048g_{2,4}g_{2,1}g_{1,4}g_{2,2} - 1536g_{1,3}g_{2,1}g_{1,4}g_{4,4} - 128g_{1,1}g_{4,4}g_{3,4}g_{3,2} - 1536g_{1,2}g_{2,3}g_{2,4}g_{3,3} + \\ & 2048g_{2,4}g_{3,3}g_{2,1}g_{1,4} - 128g_{2,2}g_{3,3}g_{1,2}g_{1,4} + 24576g_{1,3}g_{2,4}g_{3,4}g_{1,4} + 12288g_{1,4}g_{3,1}g_{2,3}g_{2,4} + 8192g_{3,4}g_{2,1}g_{2,4}g_{3,1} + 256g_{2,2}g_{3,3}g_{2,3}g_{1,4} + \\ & 512g_{3,4}g_{2,3}g_{3,1}g_{2,2} + 1792g_{3,4}g_{2,1}g_{1,1}g_{3,3} - 448g_{2,4}g_{3,1}g_{1,1}^2 + 384g_{2,3}g_{2,1}g_{1,1}g_{2,2} - 448g_{2,4}g_{3,1}g_{3,3}^2 + 1792g_{1,1}g_{3,3}g_{2,3}g_{1,4} + \\ & 4096g_{1,2}g_{2,3}g_{2,1}g_{1,4} + 2048g_{1,1}g_{4,3}g_{2,2}g_{4,4} - 448g_{2,4}g_{3,1}g_{2,2}^2 - 448g_{2,4}g_{3,1}g_{4,4}^2 + 512g_{1,1}g_{4,4}g_{1,4}g_{3,4} - 832g_{4,4}^2g_{3,4}g_{3,2} + \\ & 4096g_{1,2}g_{3,4}g_{2,1}g_{1,4} + 384g_{1,1}g_{4,4}g_{1,2}g_{1,4} + 24576g_{2,4}g_{2,1}g_{3,4} + 24576g_{2,3}g_{1,4}^2g_{3,4} + 1792g_{3,4}g_{2,1}g_{4,4}g_{2,2} + 36g_{1,1}^2g_{4,4}g_{3,3} - \\ & 256g_{4,4}g_{3,3}g_{2,3}g_{1,4} - 180g_{1,1}^2g_{4,4}g_{2,2} - 512g_{3,3}g_{2,3}g_{3,4}g_{3,1} - 1536g_{1,3}g_{1,4}^2g_{4,4} + 36g_{4,4}^2g_{2,2}g_{3,3} + 36g_{1,1}g_{3,3}^2g_{4,4} + 4096g_{2,4}g_{1,2}g_{3,1}g_{1,4} - \\ & 180g_{1,1}g_{4,4}^2g_{3,3} + 36g_{1,1}g_{4,4}^2g_{2,2} + 36g_{1,1}^2g_{2,2}g_{3,3} + 36g_{1,1}g_{2,2}^2g_{4,4} + 2048g_{4,4}g_{1,2}g_{3,4}g_{2,4} + 512g_{1,4}^2g_{1,1}g_{3,1} + 12288g_{1,4}g_{3,2}g_{3,4}^2 + \\ & 768g_{1,1}g_{2,2}g_{2,3}g_{1,4} - 2048g_{2,2}g_{2,3}g_{2,4}g_{3,4} - 512g_{1,1}g_{1,3}g_{2,1}g_{1,4} - 1536g_{3,4}^2g_{3,1}g_{4,4} - 1536g_{2,3}^2g_{2,4}g_{3,3} + 12288g_{1,3}g_{2,1}g_{2,4}g_{3,4} + \\ & 8192g_{1,2}g_{1,4}^2g_{2,3} + 12288g_{1,2}g_{1,4}^2g_{3,4} - 2048g_{1,4}g_{2,1}g_{2,4}g_{1,1} - 2048g_{2,2}g_{1,2}g_{2,4}g_{3,4} + 24576g_{2,4}g_{3,4}g_{3,1}g_{1,4} + 2048g_{1,4}g_{2,1}g_{2,4}g_{4,4} + \\ & 256g_{3,4}g_{2,1}g_{1,1}g_{2,2} + 512g_{1,1}g_{2,2}g_{3,4}g_{1,4} + 4096g_{3,4}g_{1,3}g_{3,1}g_{1,4} - 448g_{1,1}^2g_{1,3}g_{2,4} + 1536g_{1,1}g_{3,4}^2g_{3,1} - 64g_{3,3}^2g_{1,2}g_{1,4} - \\ & 512g_{1,1}g_{1,3}g_{3,1}g_{1,4}^2 + 512g_{3,4}^2g_{3,1}g_{2,2} - 180g_{1,1}g_{2,2}^2g_{3,3} + 1536g_{4,4}g_{2,3}^2g_{2,4} + 768g_{1,1}g_{4,4}g_{2,4}^2 - 512g_{2,2}g_{2,3}^2g_{2,4} + 24576g_{1,4}g_{2,1}g_{3,4}^2 + \\ & 768g_{1,1}g_{2,2}g_{2,4}^2 + 36g_{1,1}g_{2,2}g_{3,3}^2 - 180g_{4,4}g_{2,2}g_{3,3}^2 + 36g_{4,4}g_{2,2}^2g_{3,3} - 448g_{4,4}^2g_{1,3}g_{2,4} + 256g_{1,1}g_{3,3}g_{2,4}^2 + 256g_{4,4}g_{2,2}g_{2,4}^2 - \\ & 2304g_{4,4}^2g_{1,4}g_{3,4} + 12288g_{2,4}^2g_{3,4}g_{3,2} + 768g_{2,2}g_{3,3}g_{2,4}^2 + 512g_{2,3}^2g_{2,4}g_{1,1} - 2048g_{3,4}g_{2,3}g_{2,4}g_{3,3} - 448g_{2,2}^2g_{1,3}g_{2,4} + \\ & 768g_{4,4}g_{3,3}g_{2,4}^2 + 512g_{2,4}g_{2,1}^2g_{3,3} + 24576g_{2,4}^2g_{1,4}g_{2,3} + 1536g_{2,4}g_{4,4}g_{2,1}^2 - 512g_{3,3}g_{3,4}^2g_{3,1} - 448g_{1,3}g_{2,4}g_{3,3}^2 + 512g_{1,3}g_{1,4}^2g_{2,2} + \\ & 1536g_{3,3}g_{1,3}g_{1,4}^2 - 512g_{2,4}g_{2,2}g_{2,1}^2 - 1408g_{3,4}g_{2,1}g_{2,2}^2 - 832g_{4,4}^2g_{1,2}g_{1,4} - 896g_{3,4}g_{2,1}g_{1,1}^2 + 12288g_{2,4}^2g_{1,2}g_{1,4} - 1408g_{1,1}^2g_{2,3}g_{1,4} - \\ & 576g_{3,3}^2g_{3,4}g_{3,2} - 832g_{2,2}^2g_{3,4}g_{3,2} - 1408g_{2,2}^2g_{2,3}g_{1,4} - 1536g_{2,4}g_{2,1}^2g_{1,1} + 4096g_{2,4}^2g_{1,2}g_{3,2} - 2304g_{1,1}^2g_{1,4}g_{3,4} - 1408g_{3,4}g_{2,1}g_{3,3}^2 + \\ & 4096g_{2,4}^2g_{2,3}g_{3,2} - 896g_{3,3}^2g_{2,3}g_{1,4} - 1536g_{1,4}^2g_{3,3}g_{3,1} - 2304g_{3,3}^2g_{4,4}g_{1,4} - 512g_{4,4}g_{1,3}g_{3,4}^2 - 832g_{2,2}^2g_{1,2}g_{1,4} - 896g_{3,4}g_{2,1}g_{4,4}^2 - \\ & 4096g_{1,2}g_{2,3}g_{3,4}^2 - 64g_{1,1}^2g_{3,4}g_{3,2} + 1536g_{2,2}g_{1,3}g_{3,4}^2 - 896g_{4,4}^2g_{2,3}g_{1,4} + 1536g_{1,4}^2g_{3,1}g_{2,2} + 4096g_{2,4}^2g_{1,2}g_{2,1} - 2304g_{2,2}^2g_{3,4}g_{1,4} + \\ & 8192g_{2,3}^2g_{2,1}g_{1,4} - 64g_{2,3}g_{2,1}g_{4,4}^2 - 576g_{1,1}^2g_{1,2}g_{1,4} + 8192g_{3,4}g_{2,1}^2g_{2,3} + 8192g_{3,4}^2g_{3,2}g_{2,1} - 512g_{1,4}^2g_{3,1}g_{4,4} + 512g_{3,3}g_{1,3}g_{3,4}^2 - \\ & 832g_{2,3}g_{2,1}g_{1,1}^2 - 1536g_{1,1}g_{1,3}g_{3,4}^2 - 4096g_{1,4}^2g_{3,2}g_{2,1} + 12288g_{2,4}^2g_{2,1}g_{2,3} - 576g_{2,3}g_{2,1}g_{2,2}^2 - 832g_{2,3}g_{2,1}g_{3,3}^2 + 2048g_{3,4}g_{2,3}g_{2,4}g_{1,1} + \\ & 896g_{2,2}g_{3,3}g_{1,3}g_{2,4} - 128g_{4,4}g_{3,3}g_{1,3}g_{2,4} + 128g_{4,4}g_{2,2}g_{1,3}g_{2,4} + 128g_{1,1}g_{3,3}g_{1,3}g_{2,4} - 128g_{1,1}g_{2,2}g_{1,3}g_{2,4} + 896g_{1,1}g_{4,4}g_{1,3}g_{2,4} + \\ & 512g_{4,4}g_{3,3}g_{3,4}g_{1,4} + 216g_{1,1}g_{2,2}g_{4,4}g_{3,3} + 896g_{2,4}g_{3,1}g_{1,1}g_{2,2} + 896g_{2,4}g_{3,1}g_{4,4}g_{3,3} + 128g_{2,4}g_{3,1}g_{4,4}g_{2,2} + 128g_{2,4}g_{3,1}g_{1,1}g_{3,3} - \\ & 128g_{2,4}g_{3,1}g_{1,1}g_{4,4} - 128g_{2,4}g_{3,1}g_{2,2}g_{3,3} + 512g_{1,2}g_{2,3}g_{2,4}g_{1,1} - 60g_{1,1}^3g_{3,3} - 52g_{1,1}^3g_{2,2} - 52g_{1,1}^3g_{4,4} + 2048g_{2,4}^2g_{3,1}^2 - \\ & 896g_{3,3}^2g_{2,4}^2 + 150g_{4,4}^2g_{3,3}^2 + 54g_{4,4}^2g_{2,2}^2 - 52g_{4,4}^2g_{3,3}^2 - 60g_{4,4}^2g_{2,2}^2 + 54g_{1,1}^2g_{3,3}^2 + 150g_{1,1}^2g_{2,2}^2 - 52g_{1,1}g_{4,4}^3 + \\ & 150g_{1,1}^2g_{4,4}^2 - 896g_{1,1}^2g_{2,4}^2 - 8192g_{3,4}^3g_{2,3} + 8192g_{2,3}^2g_{1,4}^2 - 4096g_{1,4}^2g_{2,1}^2 - 4096g_{1,2}g_{3,4}^3 - 4096g_{2,3}^2g_{3,4}^2 - 4096g_{1,4}^3g_{3,2} + \\ & 4096g_{1,3}g_{2,4}^3 + 2048g_{1,3}^2g_{2,4}^2 - 52g_{2,2}g_{3,3}^3 + 150g_{2,2}^2g_{3,3}^2 - 896g_{2,2}^2g_{2,4}^2 - 52g_{2,2}^2g_{3,3}^2 - 8192g_{1,4}^3g_{2,1} - 52g_{4,4}g_{3,3}^3 - \\ & 60g_{4,4}g_{2,2}^3 - 896g_{4,4}^2g_{2,4}^2 - 60g_{1,1}g_{3,3}^3 - 52g_{1,1}g_{2,2}^3 + 2048g_{3,4}^2g_{3,2}^2 + 2048g_{1,2}^2g_{1,4}^2 + 8192g_{3,4}^2g_{2,1}^2 + 2048g_{2,3}^2g_{2,1}^2 + \\ & 24576g_{3,4}^2g_{1,4}^2 - 4096g_{1,4}^4 + 41g_{1,1}^4 + 41g_{4,4}^4 + 41g_{2,2}^4 + 4096g_{2,4}^3g_{3,1} - 4096g_{3,4}^4 + 41g_{3,3}^4 + 4096g_{2,4}^4 + 512g_{2,4}g_{3,2}g_{2,1}g_{3,3} \end{aligned}$$

And we have checked the following commutativity properties:

$$\{H_2, H_3\}_4 = 0,$$

$$\{H_2, H_4\}_4 = 0,$$

$$\{H_3, H_4\}_4 = 0.$$

Appendix R

Here we present the examples of the rational Lax operators for N=2,3,4. We use the following normalization:

$$L^{RT}(z) = \frac{N}{z} L(z).$$

The operator L(z) for N=2

$$L_{11} = (g_{1,1} - g_{2,2})/2$$

$$L_{12} = g_{12}$$

$$L_{21} = g_{2,1} + z^3 (g_{1,1} - g_{2,2})/2$$

$$L_{22} = -(g_{1,1} - g_{2,2})/2$$

The operator L(z) for N=3

$$L_{11} = \frac{16}{27} iz^3 g_{1,3} - 4/3 z^2 g_{2,3} - 1/3 g_{2,2} + 2/3 g_{1,1} - 1/3 g_{3,3} + 2/3 iz g_{1,2}$$

$$L_{12} = -4/3 z^2 g_{1,3} - 2 iz g_{2,3} + g_{1,2}$$

$$L_{13} = g_{1,3}$$

$$L_{21} = -\frac{32}{81} z^4 g_{1,3} - \frac{8}{27} iz^3 g_{2,3} - \frac{8}{9} z^2 g_{1,2} + (-2/3 ig_{1,1} + 2/3 ig_{3,3}) z + g_{2,1}$$

$$L_{22} = -\frac{8}{9} iz^3 g_{1,3} + 2/3 g_{2,2} - 1/3 g_{1,1} - 1/3 g_{3,3}$$

$$L_{23} = 2/3 iz g_{1,3} + g_{2,3}$$

$$L_{31} = -\frac{128}{729} z^6 g_{1,3} + \frac{32}{81} iz^5 g_{2,3} - \frac{32}{81} z^4 g_{1,2} + (-\frac{16}{27} ig_{1,1} - \frac{8}{27} ig_{3,3} + \frac{8}{9} ig_{2,2}) z^3 - 4/3 z^2 g_{2,1} - 2/3 iz g_{3,2} + g_{3,1}$$

$$L_{32} = -\frac{32}{81} iz^5 g_{1,3} - \frac{32}{27} z^4 g_{2,3} + \frac{8}{27} iz^3 g_{1,2} + (-4/3 g_{1,1} + 4/3 g_{3,3}) z^2 + 2 iz g_{2,1} + g_{3,2}$$

$$L_{33} = 4/3 z^2 g_{2,3} + 2/3 g_{3,3} - 1/3 g_{1,1} + \frac{8}{27} iz^3 g_{1,3} - 1/3 g_{2,2} - 2/3 iz g_{1,2}$$

The operator L(z) for N=4

$$L_{11} = (3/4 z^4 g_{1,4} + 2 iz^3 g_{2,4} + (-4 g_{3,4} + 3 g_{1,3}) z^2 + (4 ig_{1,2} + 4 ig_{2,3}) z - g_{3,3} - g_{4,4} + 3 g_{1,1} - g_{2,2})/4$$

$$L_{12} = iz^3 g_{1,4} - 2 z^2 g_{2,4} + (-2 ig_{3,4} + ig_{1,3}) z + g_{1,2}$$

$$L_{13} = g_{1,3} - 3/2 z^2 g_{1,4} - 2 iz g_{2,4}$$

$$L_{14} = g_{1,4}$$

$$L_{21} = \frac{3}{32} iz^5 g_{1,4} - 1/16 z^4 g_{2,4} + 3/8 iz^3 g_{1,3} + (-1/4 g_{2,3} - 3/4 g_{1,2}) z^2 + (1/2 ig_{4,4} + 1/2 ig_{3,3} - ig_{1,1}) z + g_{2,1}$$

$$L_{22} = -1/2 z^4 g_{1,4} + (-1/2 g_{1,3} - g_{3,4}) z^2 + 3/4 g_{2,2} - 1/4 g_{3,3} - 1/4 g_{4,4} - 1/4 g_{1,1}$$

$$L_{23} = -3/4 iz^3 g_{1,4} - 1/2 z^2 g_{2,4} + (1/2 ig_{1,3} - 2 ig_{3,4}) z + g_{2,3}$$

$$L_{24} = 1/2 iz g_{1,4} + g_{2,4}$$

$$L_{31} = -\frac{3}{64} z^6 g_{1,4} + 1/16 iz^5 g_{2,4} + (-1/16 g_{3,4} - 3/16 g_{1,3}) z^4 + (1/4 ig_{2,3} - 3/8 ig_{1,2}) z^3 + (3/4 g_{1,1} - 1/4 g_{3,3} - 1/2 g_{2,2}) z^2 + (-1/2 ig_{3,2} - ig_{2,1}) z + g_{3,1}$$

$$L_{32} = -1/4 iz^5 g_{1,4} - 1/2 z^4 g_{2,4} + (-1/2 ig_{3,4} - 1/4 ig_{1,3}) z^3 + (-1/4 g_{1,2} - g_{2,3}) z^2 + (-ig_{1,1} + ig_{4,4}) z + g_{3,2}$$

$$\begin{aligned}
L_{33} &= 3/8 z^4 g_{1,4} - iz^3 g_{2,4} + (-1/4 g_{1,3} + 1/2 g_{3,4}) z^2 - 1/2 iz g_{1,2} - 1/4 g_{4,4} - 1/4 g_{1,1} - 1/4 g_{2,2} + 3/4 g_{3,3} \\
L_{34} &= iz g_{2,4} - 1/4 z^2 g_{1,4} + g_{3,4} \\
L_{41} &= -\frac{3}{256} z^8 g_{1,4} + 1/16 iz^7 g_{2,4} + \left(-\frac{3}{64} g_{1,3} + \frac{3}{32} g_{3,4}\right) z^6 + \left(1/4 ig_{2,3} - \frac{3}{32} ig_{1,2}\right) z^5 + \\
&(-1/2 g_{2,2} + 3/16 g_{1,1} - 1/16 g_{4,4} + 3/8 g_{3,3}) z^4 + (-ig_{2,1} + 3/4 ig_{3,2}) z^3 \\
&+ (-1/4 g_{4,3} - 3/2 g_{3,1}) z^2 - 1/2 iz g_{4,2} + g_{4,1} \\
L_{42} &= -1/16 iz^7 g_{1,4} - 3/8 z^6 g_{2,4} + (-1/16 ig_{1,3} + 5/8 ig_{3,4}) z^5 + (-1/16 g_{1,2} - 1/2 g_{2,3}) z^4 + \\
&(-1/2 ig_{1,1} + ig_{3,3} - 1/2 ig_{4,4}) z^3 + (-1/2 g_{3,2} - 2 g_{2,1}) z^2 + (-ig_{4,3} + 2 ig_{3,1}) z + g_{4,2} \\
L_{43} &= \frac{3}{32} z^6 g_{1,4} - 5/8 iz^5 g_{2,4} + (-1/16 g_{1,3} - 5/4 g_{3,4}) z^4 + 1/2 iz^3 g_{2,3} + (-g_{2,2} + 3/2 g_{4,4} - g_{1,1} + 1/2 g_{3,3}) z^2 \\
&+ (2 ig_{2,1} + 2 ig_{3,2}) z + g_{4,3} \\
L_{44} &= -1/16 z^4 g_{1,4} + 1/2 iz^3 g_{2,4} + 3/2 z^2 g_{3,4} + (-ig_{2,3} - 1/2 ig_{1,2}) z - 1/4 g_{3,3} - 1/4 g_{2,2} + 3/4 g_{4,4} - 1/4 g_{1,1}
\end{aligned}$$

The hamiltonians and highest integrals of motion

On calculating the integrals of motion we use the standard definition:

$$H_k = \frac{1}{2k\pi i} \oint \frac{\text{tr}(L^{RT}(z))^k}{z} dz.$$

The integral H_2 is usually identified with the hamiltonian. We check the commutativity with respect to the standard Poisson bracket on the algebra $\text{gl}(N, \mathbb{C})$:

$$\{A, B\}_N = \sum_{i,j,k=1}^N \left(\frac{\partial A}{\partial g_{i,j}} \frac{\partial B}{\partial g_{j,k}} - \frac{\partial B}{\partial g_{i,j}} \frac{\partial A}{\partial g_{j,k}} \right) g_{i,k}$$

The case N=2

$$H_2 = 2 g_{1,2} (g_{1,1} - g_{2,2})$$

The case N=3

$$H_2 = -g_{1,2}^2 + 3 g_{3,3} g_{2,3} - 3 g_{2,3} g_{1,1} - 3 g_{2,1} g_{1,3}$$

$$H_3 = 3 g_{3,3} g_{2,3} g_{1,2} - 3 g_{2,3} g_{2,2} g_{1,2} + 2/3 g_{1,2}^3 - 2 g_{3,3}^2 g_{1,3} + g_{2,2}^2 g_{1,3} - 9 g_{2,3}^2 g_{2,1} + 5 g_{3,3} g_{1,1} g_{1,3} - g_{2,2} g_{1,1} g_{1,3} + 3 g_{1,3} g_{2,1} g_{1,2} - 2 g_{1,1}^2 g_{1,3} - g_{3,3} g_{2,2} g_{1,3}$$

And we check the following condition for integrability:

$$\{H_2, H_3\}_3 = 0$$

The case N=4

$$H_2 = -4 g_{1,2} g_{2,3} - 8 g_{1,4} g_{3,1} - 4 g_{2,3}^2 + 6 g_{1,1} g_{1,3} - 4 g_{2,2} g_{3,4} - 2 g_{2,2} g_{1,3} - 12 g_{1,1} g_{3,4} - 2 g_{4,4} g_{1,3} - 2 g_{3,3} g_{1,3} + 12 g_{4,4} g_{3,4} + 4 g_{3,3} g_{3,4} - 3 g_{1,2}^2 - 8 g_{2,4} g_{3,2} - 16 g_{2,4} g_{2,1}$$

$$\begin{aligned}
H_3 &= -8 g_{3,4} g_{2,1} g_{1,3} - 4 g_{1,4} g_{3,1} g_{1,2} - 4 g_{4,4} g_{3,4} g_{1,2} + 4 g_{2,4} g_{3,2} g_{1,2} - 2 g_{2,3} g_{1,3} g_{3,3} - 2 g_{1,4} g_{3,2} g_{2,2} + 4 g_{3,3} g_{2,3} g_{3,4} \\
&+ 16 g_{2,4} g_{3,1} g_{3,4} - g_{1,2}^3 - 2 g_{2,3} g_{1,3} g_{4,4} - 2 g_{1,2}^2 g_{2,3} - 2 g_{1,4} g_{3,2} g_{4,4} + 3 g_{1,1} g_{1,2} g_{1,3} - 8 g_{1,3} g_{3,1} g_{2,4} - 8 g_{1,4} g_{2,1} g_{4,4} \\
&- 4 g_{4,4} g_{2,4} g_{2,2} - 2 g_{2,3} g_{1,3} g_{2,2} - g_{2,2} g_{1,3} g_{1,2} + 4 g_{2,2} g_{2,3} g_{3,4} + 6 g_{2,3} g_{1,3} g_{1,1} + 4 g_{1,1}^2 g_{2,4} + 4 g_{4,4}^2 g_{2,4} + 16 g_{3,4}^2 g_{2,1} \\
&+ 8 g_{3,4}^2 g_{3,2} + 4 g_{4,4} g_{2,4} g_{3,3} - 4 g_{1,1} g_{2,4} g_{3,3} - 8 g_{1,1} g_{2,4} g_{4,4} - 4 g_{1,1} g_{3,4} g_{2,3} - 2 g_{1,4} g_{3,2} g_{3,3} - g_{3,3} g_{1,3} g_{1,2} + \\
&4 g_{1,1} g_{2,4} g_{2,2} + 8 g_{1,4} g_{2,1} g_{1,1} + 4 g_{2,2} g_{3,4} g_{1,2} + 6 g_{1,4} g_{3,2} g_{1,1} - g_{4,4} g_{1,3} g_{1,2} - 4 g_{4,4} g_{2,3} g_{3,4}
\end{aligned}$$

$$\begin{aligned}
H_4 &= -24 g_{2,2} g_{4,4} g_{1,2} g_{2,4} + 16 g_{1,2} g_{2,1} g_{3,4} g_{1,3} - 16 g_{2,2} g_{4,4} g_{2,4} g_{2,3} - 20 g_{2,2} g_{4,4} g_{1,1} g_{1,4} + 8 g_{1,4} g_{3,2} g_{2,3} g_{4,4} - 4 g_{2,2} g_{3,3} g_{1,1} g_{1,4} + \\
&16 g_{1,3} g_{3,1} g_{3,3} g_{1,4} + 64 g_{1,4} g_{3,2} g_{2,4} g_{3,1} + 8 g_{1,1} g_{2,2} g_{1,2} g_{2,4} - 24 g_{3,3} g_{4,4} g_{3,4} g_{1,3} - 48 g_{1,3} g_{3,2} g_{2,4} g_{1,1} + 16 g_{1,3} g_{3,2} g_{2,4} g_{4,4} + \\
&16 g_{1,4} g_{2,3} g_{2,1} g_{3,3} - 16 g_{1,1} g_{4,4} g_{1,2} g_{2,4} - 64 g_{4,4} g_{3,4} g_{1,4} g_{3,1} - 48 g_{4,4} g_{3,4} g_{1,2} g_{2,3} - 16 g_{1,4} g_{2,3} g_{2,1} g_{1,1} - 16 g_{1,4} g_{2,3} g_{2,1} g_{4,4} -
\end{aligned}$$

$$\begin{aligned}
& 256 g_{4,4} g_{3,4} g_{2,4} g_{2,1} + 16 g_{1,3} g_{3,2} g_{2,4} g_{2,2} + 16 g_{1,4} g_{2,3} g_{2,1} g_{2,2} + 12 g_{4,4} g_{1,3} g_{1,2} g_{2,3} + 80 g_{4,4} g_{1,3} g_{2,4} g_{2,1} + 16 g_{1,1} g_{3,3} g_{2,4} g_{2,3} + \\
& 8 g_{1,1} g_{3,3} g_{1,2} g_{2,4} + 16 g_{1,2} g_{2,1} g_{4,4} g_{1,4} - 32 g_{3,4} g_{1,2} g_{2,4} g_{3,1} + 32 g_{1,3} g_{3,1} g_{2,4} g_{2,3} + 16 g_{1,3} g_{3,1} g_{1,2} g_{2,4} - 48 g_{1,3} g_{3,1} g_{1,1} g_{1,4} + \\
& 16 g_{1,3} g_{3,1} g_{4,4} g_{1,4} - 64 g_{3,3} g_{3,4} g_{2,4} g_{2,1} + 12 g_{3,3} g_{1,3} g_{1,2} g_{2,3} + 16 g_{3,3} g_{1,3} g_{2,4} g_{2,1} + 16 g_{1,3} g_{3,1} g_{2,2} g_{1,4} + 16 g_{2,2} g_{1,3} g_{2,4} g_{2,1} - \\
& 8 g_{2,2} g_{4,4} g_{3,4} g_{1,3} + 8 g_{1,4} g_{3,2} g_{2,3} g_{3,3} - 16 g_{1,2} g_{2,1} g_{1,1} g_{1,4} - 8 g_{2,2} g_{3,3} g_{3,4} g_{1,3} + 4 g_{1,4} g_{3,2} g_{1,2} g_{3,3} + 4 g_{1,4} g_{3,2} g_{1,2} g_{4,4} + 56 g_{1,1} g_{4,4} g_{3,4} g_{1,3} + \\
& 56 g_{1,1} g_{3,3} g_{3,4} g_{1,3} + 8 g_{1,1} g_{2,2} g_{3,4} g_{1,3} + 8 g_{3,3} g_{4,4} g_{1,2} g_{2,4} - 12 g_{4,4}^2 g_{3,4} g_{1,3} + 48 g_{1,1} g_{3,4} g_{1,2} g_{2,3} + 256 g_{1,1} g_{3,4} g_{2,4} g_{2,1} + 64 g_{2,2} g_{3,4} g_{2,4} g_{2,1} + \\
& 20 g_{3,3} g_{4,4} g_{1,1} g_{1,4} + 4 g_{3,3} g_{4,4} g_{2,2} g_{1,4} + 32 g_{1,2} g_{2,1} g_{2,4} g_{2,3} + 16 g_{2,4} g_{3,2} g_{2,2} g_{3,4} - 48 g_{2,4} g_{3,2} g_{3,3} g_{3,4} - 112 g_{2,4} g_{3,2} g_{4,4} g_{3,4} + \\
& 144 g_{2,4} g_{3,2} g_{1,1} g_{3,4} + 12 g_{2,2} g_{1,3} g_{1,2} g_{2,3} + 64 g_{1,4} g_{3,2} g_{2,1} g_{3,4} - 8 g_{1,1}^2 g_{2,4} g_{2,3} + 32 g_{2,4} g_{2,3}^2 g_{2,1} - 32 g_{3,3} g_{2,4}^2 g_{3,1} + 40 g_{1,2}^2 g_{2,1} g_{2,4} - \\
& 32 g_{4,4} g_{2,4}^2 g_{3,1} + 10 g_{1,1}^2 g_{2,2} g_{1,4} + 8 g_{2,3}^2 g_{2,2} g_{3,4} - 8 g_{2,3}^2 g_{3,3} g_{3,4} - 56 g_{2,3}^2 g_{4,4} g_{3,4} - 2 g_{1,1}^2 g_{3,3} g_{1,4} + 64 g_{1,1} g_{3,4} g_{1,4} g_{3,1} + \\
& 28 g_{1,2}^2 g_{1,4} g_{3,1} - 56 g_{2,2} g_{4,4} g_{3,4}^2 + 56 g_{2,3}^2 g_{1,1} g_{3,4} + 4 g_{2,2}^2 g_{3,4} g_{1,3} - 21 g_{1,2}^2 g_{1,1} g_{1,3} - 104 g_{1,1} g_{4,4} g_{3,4}^2 - 8 g_{2,2} g_{3,3} g_{3,4}^2 + \\
& 2 g_{1,2}^2 g_{2,2} g_{3,4} + 7 g_{1,2}^2 g_{2,2} g_{1,3} - 10 g_{1,2}^2 g_{3,3} g_{3,4} + 7 g_{1,2}^2 g_{3,3} g_{1,3} + 40 g_{3,3} g_{4,4} g_{3,4}^2 - 26 g_{1,1}^2 g_{4,4} g_{1,4} - 12 g_{3,3}^2 g_{3,4} g_{1,3} + \\
& 128 g_{2,4} g_{2,1} g_{1,4} g_{3,1} - 36 g_{1,1} g_{1,3} g_{1,2} g_{2,3} - 112 g_{1,1} g_{1,3} g_{2,4} g_{2,1} - 2 g_{2,2}^2 g_{3,3} g_{1,4} + 6 g_{2,2}^2 g_{4,4} g_{1,4} - 64 g_{1,3} g_{3,1} g_{3,4}^2 - 12 g_{1,1} g_{3,3} g_{1,3}^2 + \\
& 64 g_{2,4}^2 g_{3,2} g_{2,1} + 64 g_{1,4} g_{2,1}^2 g_{3,4} + 12 g_{1,3} g_{2,3}^2 g_{2,2} - 36 g_{1,3} g_{2,3}^2 g_{1,1} - 60 g_{1,1}^2 g_{3,4} g_{1,3} + 32 g_{2,3} g_{3,2} g_{3,4}^2 - 12 g_{1,1} g_{4,4} g_{1,3}^2 + \\
& 4 g_{2,2} g_{4,4} g_{1,3}^2 - 12 g_{1,1} g_{2,2} g_{1,3}^2 + 4 g_{1,4} g_{3,2} g_{1,2} g_{2,2} + 32 g_{1,2} g_{2,3} g_{1,4} g_{3,1} + 4 g_{2,2} g_{3,3} g_{1,3}^2 - 56 g_{1,1} g_{3,3} g_{3,4}^2 + 4 g_{3,3} g_{4,4} g_{1,3}^2 + \\
& 40 g_{1,1} g_{2,2} g_{3,4}^2 + 30 g_{1,2}^2 g_{1,1} g_{3,4} - 8 g_{3,3}^2 g_{1,2} g_{2,4} + 8 g_{4,4}^2 g_{2,4} g_{2,3} - 12 g_{1,4} g_{3,2} g_{1,2} g_{1,1} - 6 g_{3,3}^2 g_{1,1} g_{1,4} - 8 g_{3,3}^2 g_{2,4} g_{2,3} + \\
& 2 g_{3,3}^2 g_{2,2} g_{1,4} + 16 g_{1,3} g_{3,2} g_{2,4} g_{3,3} - 2 g_{3,3}^2 g_{4,4} g_{1,4} + 2 g_{4,4}^2 g_{2,2} g_{1,4} + 12 g_{1,3} g_{2,3}^2 g_{3,3} - 10 g_{4,4}^2 g_{3,3} g_{1,4} + 16 g_{4,4}^2 g_{1,2} g_{2,4} + \\
& 26 g_{4,4}^2 g_{1,1} g_{1,4} + 8 g_{1,4} g_{3,2} g_{2,3} g_{2,2} + 16 g_{2,4} g_{2,3}^2 g_{3,2} - 24 g_{1,4} g_{3,2} g_{2,3} g_{1,1} + 32 g_{1,1} g_{2,4}^2 g_{3,1} + 32 g_{1,3} g_{2,3} g_{2,1} g_{3,4} + 32 g_{2,2} g_{2,4}^2 g_{3,1} + \\
& 32 g_{1,2} g_{2,1} g_{3,4}^2 + 12 g_{1,2}^2 g_{2,4} g_{3,2} - 22 g_{1,2}^2 g_{4,4} g_{3,4} + 7 g_{1,2}^2 g_{4,4} g_{1,3} + 12 g_{1,3} g_{2,3}^2 g_{4,4} + 48 g_{1,2} g_{3,4}^2 g_{3,2} + 16 g_{1,4} g_{2,3}^2 g_{3,1} + \\
& 8 g_{2,2}^2 g_{2,4} g_{2,3} + 8 g_{2,2}^2 g_{1,2} g_{2,4} + 2 g_{2,2}^2 g_{1,1} g_{1,4} + 4 g_{2,3}^4 + \frac{21}{4} g_{1,2}^4 + 8 g_{1,2} g_{2,3}^3 - 2 g_{2,2}^3 g_{1,4} + 12 g_{3,3}^2 g_{3,4}^2 + 60 g_{4,4}^2 g_{3,4}^2 + \\
& 14 g_{1,2}^3 g_{2,3} + 12 g_{2,2}^2 g_{3,4}^2 + 64 g_{3,4}^3 g_{3,1} + 64 g_{2,4}^2 g_{2,1}^2 + 6 g_{1,1}^3 g_{1,4} + 2 g_{3,3}^2 g_{1,3}^2 + 2 g_{4,4}^2 g_{1,3}^2 + 2 g_{2,2}^2 g_{1,3}^2 + 60 g_{1,1}^2 g_{3,4}^2 + \\
& 32 g_{1,4}^2 g_{3,1}^2 + 32 g_{2,4}^2 g_{3,2}^2 + 2 g_{3,3}^3 g_{1,4} + 18 g_{1,2}^2 g_{2,3}^2 + 18 g_{1,1}^2 g_{1,3}^2 - 6 g_{4,4}^3 g_{1,4}
\end{aligned}$$

And we check the following integrability conditions:

$$\{H_2, H_3\}_4 = 0, \quad \{H_2, H_4\}_4 = 0 \quad \{H_3, H_4\}_4 = 0$$

References

- [1] A.Smirnov, Two-Bodies systems from $SL(2, \mathbb{C})$ -tops, arXiv: math.ds/0711.2432v1 (2007)
- [2] V.I.Arnold, Mathematical Methods of Classical Mechanics, Springer, 1978 ;
- [3] M.Olshanetsky, A.Perelomov, Classical integrable finite-dimensional systems related to Lie algebras, Phys.Rep 71C 313-400 (1981)
- [4] M-P. Grosset and A.P. Veselov, Lamé equation, quantum top and elliptic Bernoulli Polynomials, arXiv: math-ph/0508068v2
- [5] A.Levin, M.Olshanetsky, A.Zotov, Hitchin Systems - Symplectic Hecke Correspondence and Two-dimensional Version. Comm.Math.Phys. 236 93-133 (2003);
- [6] N.Hitchin, Stable bundles and integrable systems, Doke Math. Jour. 54), 91-114 (1987);
- [7] A.Antonov, K.Hasegawa, A.Zabrodin, On trigonometric intertwining vectors and non-dynamical R-matrix for the Ruijsenaars model, arXiv:hep-th/9704074 (1997);
- [8] A.Gorsky, A.Zabrodin, Degenerations of Sklyanin algebra and Askey-Wilson polynomials, arXiv:hep-th/9303026 (1993).